



OFD1 gene

OFD1, centriole and centriolar satellite protein

Normal Function

The *OFD1* gene provides instructions for making a protein whose function is not fully understood. It appears to play a critical role in the early development of many parts of the body, including the brain, face, limbs, and kidneys.

The OFD1 protein is located at the base of cilia, which are finger-like projections that stick out from the surface of cells. Cilia are involved in cell movement and in many different chemical signaling pathways. They play important roles in the development and function of many parts of the body. Researchers suspect that the OFD1 protein is essential for the normal formation of cilia.

Studies suggest that the OFD1 protein may have additional functions. In the earliest stages of development, it appears to be involved in determining the left-right axis (the imaginary line that separates the left and right sides of the body). The OFD1 protein is also found in the nucleus, although its function in this cell structure is unknown.

Health Conditions Related to Genetic Changes

Joubert syndrome

oral-facial-digital syndrome

About 100 mutations in the *OFD1* gene have been found in people with oral-facial-digital syndrome type I, which is the most common form of the disorder. These mutations include changes in single DNA building blocks (base pairs) and larger deletions of genetic material from the *OFD1* gene. Most of these genetic changes lead to the production of an abnormally short, nonfunctional version of the OFD1 protein. It is unclear how a shortage of functional OFD1 protein leads to the specific features of oral-facial-digital syndrome type I. However, studies suggest that a lack of this protein prevents the normal formation of cilia, which affects the development of many tissues and organs.

primary ciliary dyskinesia

other disorders

Mutations in the *OFD1* gene can also cause several other disorders with features that overlap with those of oral-facial-digital syndrome. At least two mutations have been identified in people with Joubert syndrome. This disorder is characterized by


particular brain abnormalities, weak muscle tone (hypotonia), delayed development, unusual eye movements, and breathing problems. Another *OFD1* gene mutation has been found in a family with a form of X-linked intellectual disability. Affected individuals have had severe intellectual disability, an unusually large head size (macrocephaly), and breathing problems.

The *OFD1* mutations responsible for Joubert syndrome and X-linked intellectual disability lead to the production of an abnormally short version of the *OFD1* protein. However, studies suggest that these mutations result in a somewhat longer protein than the mutations that cause oral-facial-digital syndrome. This protein may retain some of its usual function in cilia. The abnormal protein probably disrupts the development or function of cilia in some way, although it is unclear how these changes result in the signs and symptoms of Joubert syndrome or X-linked intellectual disability.

Chromosomal Location

Cytogenetic Location: Xp22.2, which is the short (p) arm of the X chromosome at position 22.2

Molecular Location: base pairs 13,734,713 to 13,769,361 on the X chromosome (Homo sapiens Annotation Release 108, GRCh38.p7) (NCBI)



Credit: Genome Decoration Page/NCBI

Other Names for This Gene

- 71-7A
- CXorf5
- JBTS10
- MGC117039
- MGC117040
- OFD1_HUMAN
- oral-facial-digital syndrome 1
- SGBS2

Additional Information & Resources

Educational Resources

- Kennedy Krieger Institute: X-linked Mental Retardation
<http://gcbcenter.kennedykrieger.org/xmr.jsp>
- Molecular Biology of the Cell (fourth edition, 2002): Cilia and Flagella Are Motile Structures Built from Microtubules and Dyneins
<https://www.ncbi.nlm.nih.gov/books/NBK26888/#A3075>
- Molecular Cell Biology (first edition, 2000): Cilia and Flagella: Structure and Movement
<https://www.ncbi.nlm.nih.gov/books/NBK21698/>

GeneReviews

- Joubert Syndrome and Related Disorders
<https://www.ncbi.nlm.nih.gov/books/NBK1325>
- Oral-Facial-Digital Syndrome Type I
<https://www.ncbi.nlm.nih.gov/books/NBK1188>

Scientific Articles on PubMed

- PubMed
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28OFD1%5BTIAB%5D%29+OR+%28%28SGBS2%5BTIAB%5D%29+OR+%28CXorf5%5BTIAB%5D%29%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+3600+days%22%5Bdp%5D>

OMIM

- JOUBERT SYNDROME 10
<http://omim.org/entry/300804>
- OFD1 GENE
<http://omim.org/entry/300170>

Research Resources

- ClinVar
<https://www.ncbi.nlm.nih.gov/clinvar?term=OFD1%5Bgene%5D>
- HGNC Gene Symbol Report
http://www.genenames.org/cgi-bin/gene_symbol_report?q=data/hgnc_data.php&hgnc_id=2567

- NCBI Gene
<https://www.ncbi.nlm.nih.gov/gene/8481>
- UniProt
<http://www.uniprot.org/uniprot/O75665>

Sources for This Summary

- Budny B, Chen W, Omran H, Fliegauf M, Tzschach A, Wisniewska M, Jensen LR, Raynaud M, Shoichet SA, Badura M, Lenzner S, Latos-Bielecka A, Ropers HH. A novel X-linked recessive mental retardation syndrome comprising macrocephaly and ciliary dysfunction is allelic to oral-facial-digital type I syndrome. *Hum Genet.* 2006 Sep;120(2):171-8. Epub 2006 Jun 17.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/16783569>
- Coene KL, Roepman R, Doherty D, Afroze B, Kroes HY, Letteboer SJ, Ngu LH, Budny B, van Wijk E, Gorden NT, Azhimi M, Thauvin-Robinet C, Veltman JA, Boink M, Kleefstra T, Cremers FP, van Bokhoven H, de Brouwer AP. OFD1 is mutated in X-linked Joubert syndrome and interacts with LCA5-encoded lebercillin. *Am J Hum Genet.* 2009 Oct;85(4):465-81. doi: 10.1016/j.ajhg.2009.09.002.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/19800048>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2756557/>
- Ferrante MI, Giorgio G, Feather SA, Bulfone A, Wright V, Ghiani M, Selicorni A, Gammaro L, Scolari F, Woolf AS, Sylvie O, Bernard L, Malcolm S, Winter R, Ballabio A, Franco B. Identification of the gene for oral-facial-digital type I syndrome. *Am J Hum Genet.* 2001 Mar;68(3):569-76. Epub 2001 Feb 13.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/11179005>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1274470/>
- Ferrante MI, Zullo A, Barra A, Bimonte S, Messaddeq N, Studer M, Dollé P, Franco B. Oral-facial-digital type I protein is required for primary cilia formation and left-right axis specification. *Nat Genet.* 2006 Jan;38(1):112-7. Epub 2005 Nov 27.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/16311594>
- Giorgio G, Alfieri M, Prattichizzo C, Zullo A, Cairo S, Franco B. Functional characterization of the OFD1 protein reveals a nuclear localization and physical interaction with subunits of a chromatin remodeling complex. *Mol Biol Cell.* 2007 Nov;18(11):4397-404. Epub 2007 Aug 29.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/17761535>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2043566/>
- Macca M, Franco B. The molecular basis of oral-facial-digital syndrome, type 1. *Am J Med Genet C Semin Med Genet.* 2009 Nov 15;151C(4):318-25. doi: 10.1002/ajmg.c.30224. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/19876934>
- Prattichizzo C, Macca M, Novelli V, Giorgio G, Barra A, Franco B; Oral-Facial-Digital Type I (OFDI) Collaborative Group. Mutational spectrum of the oral-facial-digital type I syndrome: a study on a large collection of patients. *Hum Mutat.* 2008 Oct;29(10):1237-46. doi: 10.1002/humu.20792.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/18546297>
- Romio L, Fry AM, Winyard PJ, Malcolm S, Woolf AS, Feather SA. OFD1 is a centrosomal/basal body protein expressed during mesenchymal-epithelial transition in human nephrogenesis. *J Am Soc Nephrol.* 2004 Oct;15(10):2556-68.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/15466260>

- Thauvin-Robinet C, Cossée M, Cormier-Daire V, Van Maldergem L, Toutain A, Alembik Y, Bieth E, Layet V, Parent P, David A, Goldenberg A, Mortier G, Héron D, Sagot P, Bouvier AM, Huet F, Cusin V, Donzel A, Devys D, Teyssier JR, Faivre L. Clinical, molecular, and genotype-phenotype correlation studies from 25 cases of oral-facial-digital syndrome type 1: a French and Belgian collaborative study. *J Med Genet.* 2006 Jan;43(1):54-61.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/16397067>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2564504/>
 - Thauvin-Robinet C, Franco B, Saugier-Veber P, Aral B, Gigot N, Donzel A, Van Maldergem L, Bieth E, Layet V, Mathieu M, Teebi A, Lespinasse J, Callier P, Mugneret F, Masurel-Paulet A, Gautier E, Huet F, Teyssier JR, Tosi M, Frébourg T, Faivre L. Genomic deletions of OFD1 account for 23% of oral-facial-digital type 1 syndrome after negative DNA sequencing. *Hum Mutat.* 2009 Feb;30(2):E320-9. doi: 10.1002/humu.20888.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/19023858>
-

Reprinted from Genetics Home Reference:

<https://ghr.nlm.nih.gov/gene/OFD1>

Reviewed: February 2010

Published: March 21, 2017

Lister Hill National Center for Biomedical Communications
U.S. National Library of Medicine
National Institutes of Health
Department of Health & Human Services